

Observing the diffuse supernova neutrino background



SN 1987A, Anglo-Australian
Observatory/David Malin Images

Peter Madigan

Outline


What is the diffuse supernova neutrino background (DSNB)?

Why search for the DSNB?

Recent DSNB searches

Future of the DSNB

The lifecycle of a star




Stars fuse light nuclei into heavier and heavier nuclei.

Requiring hotter temperatures to fuse.

Iron ends the fusion cycle.


The lifecycle of a star



Stars fuse light nuclei into heavier and heavier nuclei.

Requiring hotter temperatures to fuse.

Iron ends the fusion cycle.




The inward gravitational pressure of the core eventually overcomes the outward thermal/e-degeneracy pressure. ($>8M$)

Collapsing the core into neutron star.

Neutrino emission

During collapse,



Core is on the order of nuclear densities so the neutrino scattering length is appreciable:


Most of the energy is released through neutrinos.

Neutrinos are likely emitted with a thermal spectrum.

$$\phi(E_\nu) = E_{\bar{\nu}_e, tot} \frac{120}{7\pi^4} \frac{E_\nu^2}{T^4} \frac{1}{e^{E_\nu/T} + 1}$$

Neutrino emission

During collapse,



Core is on the order of nuclear densities so the neutrino scattering length is appreciable:

Most of the energy is released through neutrinos.


Neutrinos are likely emitted with a thermal spectrum.

$$\phi(E_\nu) = E_{\bar{\nu}_e, tot} \frac{120}{7\pi^4} \frac{E_\nu^2}{T^4} \frac{1}{e^{E_\nu/T} + 1}$$


But how likely are we to see one of these?

Supernova rate

Not *that* likely... about 1 supernova within the Milky Way every 20-50 years.
Last one in 1987:



K. Hirata *et al.*, "Observation of a neutrino burst from the supernova SN1987A," *Phys. Rev. Lett.*, vol. 58, no. 14, pp. 1490–1493, Apr. 1987.



SN 1987A, NASA 2007.

So do we just hope for another one in our lifetime?


“Space is big. Really big. You just won't believe how vastly, hugely, mind-bogglingly big it is. I mean, you may think it's a long way down the road to the chemist, but that's just peanuts to space.”

-Douglas Adams

At any given moment, there should be *some* neutrinos reaching Earth from some distant supernova.

A number density using the supernova rate as a function of redshift:

$$\frac{dn_\nu}{dE_\nu} = \int R_{SN}(z)(1+z)\phi(E'_\nu)\frac{dt}{dz}dz$$



S. Ando and K. Sato, “Relic neutrino background from cosmological supernovae,” *New Journal of Physics*, vol. 6, pp. 170–170, Nov. 2004. 8

Why look for diffuse supernova neutrinos?

For astrophysics:

- DSNB measurements could be used to find star-formation rates and supernova rates, unaffected by interstellar dust.
- SN come directly from the core of the collapsing star and are the most sensitive probe of the physics that occurs in this process.

For particle physics:

- Flavor make-up of the DSNB is sensitive to the neutrino mass hierarchy and mixing angles.
- The long-baseline of the DSNB is sensitive to neutrino decay, which would have broad implications in particle physics and in astrophysics.


Looking for the DSNB:

Roughly equal portions of all neutrino flavors.

Thermal spectrum peaked at about 4-8 MeV.


Isotropic.

Flux comparable to low-energy atmospheric neutrinos.




Low energy excludes CC interactions for muon and tau neutrinos.
Cross-sections make NC/elastic scattering unlikely.


Observation will likely be made through an inverse beta decay search.



Signal



Signal



Signal


Generic detector

- Cherenkov light
- Scintillation
- Ionization

[1] K. Bays *et al.*, “Supernova relic neutrino search at Super-Kamiokande,” *Physical Review D*, vol. 85, no. 5, Mar. 2012.

[2] H. Zhang *et al.*, “Supernova Relic Neutrino search with neutron tagging at Super-Kamiokande-IV,” *Astroparticle Physics*, vol. 60, pp. 41–46, Jan. 2015.

[3] A. Gando *et al.*, “Search for extraterrestrial antineutrino sources with the KamLAND detector,” *The Astrophysical Journal*, vol. 745, no. 2, p. 193, Feb. 2012.



- Pair-production
- Scintillation
- Ionization

Super-Kamiokande



50kt water Cherenkov detector buried 1000m underground in the Kamioka mine (Japan).

Operating since 1996, published bounds on the DSNB in 2003, 2012, 2015, using two different methods:



- Only positron events
- Positron with neutron tagging

Biggest backgrounds:

- Invisible-muon decays (higher energy)
- NC elastic scattering (lower energy)



Super-Kamiokande






KamLAND



1kt liquid scintillator detector also in Kamioka mine. Running from 2002-11, searches for the DSNB through delayed coincidence.

Backgrounds:

- Spallation with positron and neutron in f.s.
- NC interactions with nuclei




KamLAND






But can we do better..?

Of course! Super-Kamiokande has an inverse beta decay efficiency of only 13%.



GADZOOKS! (Gadolinium Antineutrino Detector Zealously Outperforming Old Kamiokande, Super!)



J. F. Beacom and M. R. Vagins, "Antineutrino Spectroscopy with Large Water Cerenkov Detectors," *Phys. Rev. Lett.*, vol. 93, no. 17, p. 171101, Oct. 2004.



Figures: P. Fernandez, "Status of GADZOOKS!: Neutron Tagging in Super-Kamiokande," in *Nuclear Physics B Proceedings Supplement 00* (2014), pp. 1–8.

Others (more distant future)


Liquid argon detector (DUNE):




www.dunescience.org



Water-based liquid scintillator (ASDC/THEIA):




arxiv:1504.08284




arxiv:1409.5864

Large liquid scintillator detector (JUNO):



arxiv:1507.05613

290E - Peter Madigan



Last thoughts

- Supernova neutrinos are useful to both astrophysics and neutrino physics.
- The DSNB gives neutrino experiments something to strive for while also preparing for the next near-by supernova.
- In recent history, we have been able to get close to observing the DSNB (likely within a factor of <10).
- The DSNB is observable in the near future!

References

- [1] K. Bays *et al.*, “Supernova relic neutrino search at Super-Kamiokande,” *Physical Review D*, vol. 85, no. 5, Mar. 2012.
 - [2] H. Zhang *et al.*, “Supernova Relic Neutrino search with neutron tagging at Super-Kamiokande-IV,” *Astroparticle Physics*, vol. 60, pp. 41–46, Jan. 2015.
 - [3] A. Gando *et al.*, “Search for extraterrestrial antineutrino sources with the KamLAND detector,” *The Astrophysical Journal*, vol. 745, no. 2, p. 193, Feb. 2012.
 - [4] S. Horiuchi, J. F. Beacom, and E. Dwek, “Diffuse supernova neutrino background is detectable in Super-Kamiokande,” *Phys. Rev. D*, vol. 79, no. 8, p. 083013, Apr. 2009.
- + others where cited.